

Manifestation of a nontrivial vacuum in discrete light cone quantization

Takanori Sugihara and Masa-aki Taniguchi

Department of Physics, Nagoya University, Chikusa, Nagoya 464-8602, Japan

We study a (1+1)-dimensional $\lambda\phi^4$ model with a light-cone zero mode and constant external source to describe spontaneous symmetry breaking. In the broken phase, we find degenerate vacua and discuss their stability based on effective-potential analysis. The vacuum triviality is spurious in the broken phase because these states have lower energy than Fock vacuum. Our results are based on the variational principle.

Light-cone quantization [1] has been studied to clarify nonperturbative aspects of field theories [2] and used to provide nonperturbative formulation of M-theory [3]. This framework simplifies dynamics of quantum field theories since it prohibits vacuum diagrams kinematically [4]. It has also a possibility of calculating wavefunctions of physical states in a nonperturbative manner. Because of these preferable properties, we hope for this framework playing a complementary role to lattice theories.

DLCQ (discrete light cone quantization) is the most well-defined treatment of light-cone quantization, which enables clear separation of zero mode [5–7]. It has been believed that the true vacuum is trivial and only the zero mode is responsible for spontaneous symmetry breaking (SSB) in scalar field theories [5,8]. There are some studies that rely on a combination of constrained zero modes and trivial vacuum [9]. The vacuum triviality, however, results from an assumption that normal-ordered Hamiltonians are positive semidefinite. In this letter, we show that this assumption is not always true. We include a zero mode and external source in analytic variational calculations to show the existence of nontrivial vacuum with lower energy than trivial Fock vacuum. Our results are based on previous works on zero-mode singularity [8] and quantum solitons [10,11] in DLCQ. We examine a possibility that zero modes reproduce SSB, which has not been considered in Ref. [10,11].

In order to define an effective potential, we consider the generating functional of Green's functions, $W[J] = -i \ln Z[J]$. We can express it using a Hamiltonian $H[J]$ when the external source is time-independent $J(x) = J(\mathbf{x})$ [10,12],

$$H[J]|0_J\rangle = w[J]|0_J\rangle, \quad H[J] = H - \int_{-L}^L d\mathbf{x} J(\mathbf{x})\phi(\mathbf{x}),$$

where $w[J] = -W[J]/T$ is energy of the ground state $|0_J\rangle$. T is the time difference between the initial and final states. If we are just interested in the ground state (vacuum), we can consider an effective potential to reduce the problem into simpler one. It is defined as a Legendre transform $V(\varphi_0) \equiv w(J) + J\varphi_0 = \langle 0_J | H | 0_J \rangle / 2L$ with a constant source $J(x) = J$ and vacuum-energy density $w(J) = w[J]/2L$. A vacuum expectation value (VEV) of a zero mode is given by $\varphi_0 = -dw(J)/dJ = \langle 0_J | \phi_0 | 0_J \rangle$. We can calculate the effective potential $V(\varphi_0)$ if energy and wavefunction of the ground state are known. Our purpose is to identify the true vacuum in the (1+1)-dimensional $\lambda\phi^4$ model with a double-well classical potential. We perform variational calculations for the following normal-ordered Hamiltonian.

Hereafter, we will designate the space coordinate x^- by x . The field operator is decomposed into zero and nonzero modes $\phi(x) = \phi_0 + \tilde{\phi}(x)$, where $(O)_0 \equiv \int_{-L}^L dx O(x) / 2L$. In DLCQ with periodic boundary conditions, the zero mode ϕ_0 is constrained [5],

$$H(J) = \int_{-L}^L dx^- \left(-\frac{\mu^2}{2} : \phi^2 : + \frac{\lambda}{4!} : \phi^4 : - J : \phi : \right). \quad (1)$$

Operator ordering has been chosen so that it satisfies $dw(J)/dJ = -\varphi_0$. It is the consistency condition to be satisfied between the Hamiltonian (1) and zero-mode constraint (2). We will discuss it later in Eq. (24). The zero mode is an operator functional of the nonzero mode

$$\Phi[\tilde{\phi}] = -\mu^2 : \phi_0 : + \frac{\lambda}{6} : (\phi^3)_0 : - J = 0. \quad (2)$$

where $[a_m, a_n^\dagger] = \delta_{m,n}$ and $a_m|0_F\rangle = 0$. We truncate the system by introducing one-coherent state $|\tilde{\varphi}\rangle = e^{\sum_{n=1}^{\infty} (\tilde{\varphi}_n a_n^\dagger - \frac{1}{2} \tilde{\varphi}_n^* \tilde{\varphi}_n)} |0_F\rangle$. It provides useful formulas $a_n|\tilde{\varphi}\rangle = \tilde{\varphi}_n|\tilde{\varphi}\rangle$ and $\langle \tilde{\varphi} | : O[\tilde{\phi}] : | \tilde{\varphi} \rangle = O[\tilde{\varphi}]$, where $\varphi(x) \equiv \langle \tilde{\varphi} | \phi(x) | \tilde{\varphi} \rangle = \varphi_0 + \tilde{\varphi}(x)$. Our variational parameter is $\tilde{\varphi}(x)$.

We take the continuum limit with $P^+ = \pi M/L$ fixed, where M is called the harmonic resolution. We include this constraint on P^+ in Hamiltonian using the Lagrange's undetermined multiplier.

We take the continuum limit with $P^+ = \pi M/L$ fixed, where M is called the harmonic resolution. We include this constraint on P^+ in Hamiltonian using the Lagrange's undetermined multiplier.

We take the continuum limit with $P^+ = \pi M/L$ fixed, where M is called the harmonic resolution. We include this constraint on P^+ in Hamiltonian using the Lagrange's undetermined multiplier.

$$E \equiv \langle \tilde{\varphi} | H_\beta(J) | \tilde{\varphi} \rangle \quad (4)$$

$$= \int_{-L}^L dx \left[\beta \left(\left(\frac{d\tilde{\varphi}}{dx} \right)^2 - \frac{\pi M}{2L^2} \right) - \frac{\mu^2}{2} \varphi^2 + \frac{\lambda}{4!} \varphi^4 - J\varphi \right],$$

where $w(J) = E/2L$. Equation (2) and the stationary condition for (4) give the following coupled equations for the zero and nonzero modes.

$$-\mu^2\varphi_0 + \frac{\lambda}{6}\varphi_0^3 = J, \quad (5)$$

$$-2\beta\frac{d^2\tilde{\varphi}}{dx^2} + \mu_e^2\tilde{\varphi} + \frac{\lambda}{2}\varphi_0\tilde{\varphi}^2 + \frac{\lambda}{6}\tilde{\varphi}^3 = 0, \quad (6)$$

where $\mu_e^2 \equiv -\mu^2 + \lambda\varphi_0^2/2$. We have the following zero-mode solutions to (5) when $J = 0$ (see Fig. 1).

$$\varphi_0 = 0, \pm v, \quad v \equiv \sqrt{\frac{6\mu^2}{\lambda}}. \quad (7)$$

The system defined in a finite box can describe SSB if the vanishing J limit is taken after all calculations. We should choose $\varphi_0 = \pm v$ but not $\varphi_0 = 0$ as physical solutions to (5) when $J = 0$, because the solution $\varphi_0 = 0$ is not connected continuously to expectation values for large $J \neq 0$. The region $|\varphi_0| < v/\sqrt{3}$ is unphysical.




FIG. 1. Functional relationship (5) between φ_0 and J is shown when Fock space is truncated with the one-coherent approximation. The region $|\varphi_0| < v/\sqrt{3}$ is unstable (see discussions given below Eq. (25)). When $|\varphi_0| \geq v$, zero mode φ_0 increases monotonously, $d\varphi_0/dJ > 0$.

If one attempts to see vacuum physics without introducing an external source, wrong solutions may be obtained. Symmetric phase would be safe without an external source, but broken phase is not. We will explain how the true vacuum solution appears by paying attention to the effects of the zero mode and external source on the ground state. We will also discuss vacuum stability in Eq. (25) based on the effective potential.

Solutions to Eq. (6) must satisfy the following two conditions simultaneously:

- (i) The solution $\tilde{\varphi}$ to (6) and the n -th derivative $\tilde{\varphi}^{(n)} = d^n\tilde{\varphi}/dx^n$ must be periodic at the boundaries $x = \pm L$: $\tilde{\varphi}^{(n)}(-L) = \tilde{\varphi}^{(n)}(L)$, $n = 0, 1, 2, \dots$

- (ii) The solution $\tilde{\varphi}$ to (6) must be nonzero mode: $(\tilde{\varphi}(x))_0 = \int_{-L}^L dx \tilde{\varphi}(x)/2L = 0$.

For convenience, let us regard x as time and consider the following dummy Hamiltonian $\mathcal{H}$ that reproduces the equation of motion (6).

$$\mathcal{H} = \frac{1}{2}(\partial\tilde{\varphi})^2 + \mathcal{V}(\tilde{\varphi}), \quad (8)$$

where $\mathcal{V}$ is a dummy potential

$$\mathcal{V}(\tilde{\varphi}) = -\frac{1}{2\beta} \left(\frac{\mu_e^2}{2}\tilde{\varphi}^2 + \frac{\lambda}{6}\varphi_0\tilde{\varphi}^3 + \frac{\lambda}{4!}\tilde{\varphi}^4 - \frac{\mu^2}{2}\varphi_0^2 + \frac{\lambda}{4!}\varphi_0^4 \right). \quad (9)$$

We first solve (6) especially when $J = 0$ and $\varphi_0 = 0$. As mentioned before, this gives physically unacceptable solutions. However, these solutions are technically helpful for the purpose of calculating energy and wavefunction of the true degenerate vacua with nonzero VEVs as shown later. When $J = 0$ and $\varphi_0 = 0$, Eq. (6) reduces to

$$\frac{d^2\tilde{\varphi}}{dx^2} = \frac{1}{2\beta} \left(-\mu^2\tilde{\varphi} + \frac{\lambda}{6}\tilde{\varphi}^3 \right), \quad (10)$$

and the dummy potential $\mathcal{V}$ is

$$\mathcal{V}(\tilde{\varphi}) = -\frac{1}{2\beta} \left(-\frac{\mu^2}{2}\tilde{\varphi}^2 + \frac{\lambda}{4!}\tilde{\varphi}^4 \right). \quad (11)$$

When the parameter β is positive, the dummy potential $\mathcal{V}(\tilde{\varphi})$ is bounded from above (see Fig. 2(a)). From the condition (i), motion of a particle must be periodic. Namely, the particle must reside between the two maximums of the dummy potential. If the condition (i) is satisfied, the condition (ii) is also satisfied since the particle oscillates around the origin in the symmetric dummy potential. The solution to (10) is

$$\tilde{\varphi}_{\text{sn}}(x) = \left[\frac{12\mu^2 k^2}{\lambda(k^2 + 1)} \right]^{1/2} \text{sn}(a_{\text{sn}}x, k), \quad (12)$$

where sn is a Jacobian elliptic function and $0 \leq k \leq 1$ [11]. The values of the parameters a_{sn} and k are determined so that the solution (12) satisfies both the conditions (i) and (ii); $k = 0$ gives $\tilde{\varphi}(x) = 0$ and $M = 0$, which correspond to trivial Fock vacuum with $\varphi_0 = 0$ and $P^- = 0$. $k = 1$ is not acceptable since it gives $\tilde{\varphi} \sim \tanh(ax)$, which is an odd function and cannot satisfy the periodicity condition (i). When $0 < k < 1$, we have $a_{\text{sn}} = 2NK(k)/L$ (N is a natural number and $K(k)$ is the complete elliptic integral of the first kind) from the periodicity condition (i) and it satisfies the condition (ii). In this case, the solution (12) is acceptable and gives the following energy and harmonic resolution.

$$\frac{E_{\text{sn}}}{2L} = -\frac{6\mu^4 k^2}{\lambda(k^2 + 1)^2} + \frac{576N^2\mu^6 k^4 I_{\text{sn}}^2(k)}{\pi\lambda^2(k^2 + 1)^3 M}, \quad (13)$$

$$M = \frac{96N\mu^2 k^2 I_{\text{sn}}(k)K(k)}{\pi\lambda(k^2 + 1)}, \quad (14)$$

$$I_{\text{sn}}(k) \equiv \int_0^1 df \sqrt{(1-f^2)(1-k^2 f^2)}. \quad (15)$$

In the limit $k \rightarrow 1$ with $N = 1$, the energy (13) takes the minimum value, and the harmonic resolution goes to infinity since $K(k)$ diverges at $k = 1$ giving the continuum limit $M \rightarrow \infty$. This is the solution given in Ref. [11].




FIG. 2. The dummy potential $\mathcal{V}$ is drawn as a function of $\tilde{\varphi}$ when $\varphi_0 = 0$: (a) positive β and (b) negative β .

When the parameter β is negative, the condition (i) is automatically satisfied since the dummy potential $\mathcal{V}(\tilde{\varphi})$ is bounded from below and a particle oscillates necessarily with a fixed period (see Fig. 2(b)). Equation (10) has two types of solutions in this case. We can also express them using Jacobian elliptic functions $\text{cn}(x, k)$ and $\text{dn}(x, k)$. When a particle oscillates around the origin, a solution is

$$\tilde{\varphi}_{\text{cn}}(x) = \left[\frac{12\mu^2 k^2}{\lambda(2k^2 - 1)} \right]^{1/2} \text{cn}(a_{\text{cn}} x, k). \quad (16)$$

The solution (16) is acceptable as an excited state since it satisfies the conditions (i) and (ii) simultaneously with higher energy than (12). When a particle oscillates around one of the minimums of the dummy potential, solutions are

$$\tilde{\varphi}_{\text{dn}}(x) = \pm \left[\frac{12\mu^2}{\lambda(2 - k^2)} \right]^{1/2} \text{dn}(a_{\text{dn}} x, k). \quad (17)$$

The solutions (17) are not acceptable since they cannot satisfy the condition (ii).

When $J = 0$ and $\varphi_0 = 0$, the candidate solution for the ground state is (12) with $a_{\text{sn}} = 2NK(k)/L$ and $k \rightarrow 1$ derived for positive β since it gives the lowest energy. However, we discard the solution (12) since the state formed by it cannot be connected to solutions for non-zero external source $J \neq 0$. In addition, mass squared goes to negative infinity $P^\mu P_\mu \rightarrow -\infty$ in the continuum limit $M \rightarrow \infty$ since the first term of the energy (13) is nonzero and negative. This is the reason why all past calculations have not been successful in calculating mass squared stably in the broken phase [10].

When the limit $J \rightarrow 0$ is taken starting from a sufficiently large J , an effective potential chooses one of $\varphi_0 = \pm v$ depending on the sign of J . Nonzero values of J resolve the degeneracy of the two vacua, which restores in the limit $J \rightarrow 0$. In this case, the conditions (i) and (ii) select dn-type oscillation around the minimum $\tilde{\varphi} = 0$ of the dummy potential. The situation is completely different from the $\varphi_0 = 0$ case. We obtain the following dummy potential by substituting $\varphi_0 = v$ into (9) (it is enough to consider one of the two degenerate vacua $\varphi_0 = \pm v$).

$$\mathcal{V}(\tilde{\varphi}) = -\frac{1}{2\beta} \left(\mu^2 \tilde{\varphi}^2 + \frac{\lambda}{6} v \tilde{\varphi}^3 + \frac{\lambda}{4!} \tilde{\varphi}^4 - \frac{3\mu^4}{2\lambda} \right). \quad (18)$$

This is the case when the nonzero mode is shifted with $\tilde{\varphi} \rightarrow \tilde{\varphi} + v$ in (11). Therefore, we obtain four possible oscillations by shifting the solutions (12), (16), and (17) with $\tilde{\varphi} \rightarrow \tilde{\varphi} - v$. The first is sn-type oscillation around $\tilde{\varphi} = -v$. The second is cn-type oscillation around $\tilde{\varphi} = -v$. The third is dn-type oscillation around $\tilde{\varphi} = -2v$. However, all of them are unacceptable since they cannot satisfy the condition (ii). The following dn-type solution is a physically acceptable oscillation:

$$\tilde{\varphi}(x) = \left[\frac{12\mu^2}{\lambda(2 - k^2)} \right]^{1/2} \text{dn}(ax, k) - v, \quad (19)$$

which oscillates around the origin $\tilde{\varphi} = 0$ and can satisfy the conditions (i) and (ii) simultaneously. Its energy and total momentum are

$$\frac{E}{2L} = \frac{6\mu^4(k^2 - 1)}{\lambda(2 - k^2)^2} - \frac{144\mu^6 I_{\text{dn}}^2(k)}{\pi\lambda^2(2 - k^2)^3 M}, \quad (20)$$

$$P^+ = \frac{\pi M}{L} = \frac{24\mu^2 I_{\text{dn}}(k)}{\lambda(2 - k^2)} a, \quad (21)$$

$$I_{\text{dn}}(k) \equiv \int_{\text{dn}(aL, k)}^1 df \sqrt{(1 - f^2)(f^2 - 1 + k^2)},$$

where $0 \leq k \leq 1$; $k = 0$ is not acceptable since $\tilde{\varphi}(x) = 0$ gives $\varphi_0 = \langle 0_{\text{F}} | : \phi_0 : | 0_{\text{F}} \rangle = 0$ that contradicts $\varphi_0 = v$. When $0 < k < 1$, there exists no value of the parameter $a = NK(k)/L$ that can satisfy the condition (ii). When $k = 1$, we have

$$\tilde{\varphi}(x) = v \left[\frac{\sqrt{2}}{\cosh(ax)} - 1 \right]. \quad (22)$$

The condition (ii) requires the following relation to hold.

$$\sqrt{2} \text{gd}(aL) - aL = 0, \quad (23)$$

where gd is the Gudermann function. This has a solution $aL \sim 1.72$. In the continuum limit $L \rightarrow \infty$, we have $a \rightarrow 0$ which gives zero total momentum $P^+ = 0$. Since the energy (20) is negative and lower than Fock vacuum

in the continuum limit, we identify it as one of the true vacua.

In order to check the consistency of the operator ordering between Hamiltonian (1) and zero-mode constraint (2), we examine the first derivative of $w(J)$ with respect to J using (5) and (6).

$$\begin{aligned} & \frac{dw}{dJ} + \varphi_0 \\ &= \int_{-L}^L \frac{dx}{2L} \left[\frac{d\tilde{\varphi}}{dJ} \left(\mu_e^2 \tilde{\varphi} + \frac{\lambda}{2} \varphi_0 \tilde{\varphi}^2 + \frac{\lambda}{6} \tilde{\varphi}^3 \right) + \frac{d\varphi_0}{dJ} \Phi[\tilde{\varphi}] \right] \\ &= -\beta \frac{d}{dJ} \left(\frac{P^+}{2L} \right). \end{aligned} \quad (24)$$

Since the parameters P^+ and L are given by hand independent of J , we obtain the desired consistency condition $dw/dJ = -\varphi_0$. We can use this relation to discuss vacuum stability also. We have the following relation from the definition of the effective potential $V(\varphi_0) = w(J) + J\varphi_0$.

$$\frac{d^2 V}{d\varphi_0^2} = \frac{dJ}{d\varphi_0}. \quad (25)$$

The state given by (22) is stable and hence can be regarded as one of the true vacua since (25) is positive when $\varphi_0 = v$. On the other hand, the state given by (12) is unstable since (25) is negative when $\varphi_0 = 0$. In $-v/\sqrt{3} < \varphi_0 < v/\sqrt{3}$, energy decreases as J increases. If ϕ_0 or J is not introduced, one cannot observe this instability of the state (12) since $V(\varphi_0)$ and hence (25) are not available.

We conclude that there exist nontrivial degenerate vacua other than Fock vacuum in the (1+1)-dimensional $\lambda\phi^4$ model with a double-well classical potential. We have shown stability of the obtained vacua based on the effective-potential analysis. The essential point of our analysis is introduction of a zero mode and external source.

In general, there are singlet and non-singlet sectors of Z_2 symmetry. One-coherent state $|\tilde{\varphi}\rangle$ is a mixed state of both the sectors. We have shown that there exist non-singlet vacua with lower energy than Fock vacuum when the classical potential has a double-well shape. The mixing of the singlet and non-singlet sectors is a consequence of introduction of an explicitly symmetry-breaking interaction $J\phi_0$ in the Hamiltonian (1).

The issues of critical exponents remain still open until quantitatively reliable calculations are done. We should perform variational calculations without assuming vacuum triviality also in the case when a classical potential is convex. However, we will need to interpret the difference among normal (ours) and other operator orderings (such as Weyl ordering) concerning to the origin of VEVs of zero modes. When the Hamiltonian and zero-mode constraint are normal-ordered, trivial Fock vacuum

just gives $\langle 0_F | : H : | 0_F \rangle = 0$ and $\langle 0_F | : \phi_0 : | 0_F \rangle = 0$ (i.e. there is no SSB if vacuum triviality is assumed for normal-ordered operators).

Finally, we point out the importance of small momentum components near the zero mode. The reason why the true-vacuum solution (22) gives $P^+ = 0$ in the continuum limit $L \rightarrow \infty$ is that its slowly-changing configuration is mainly composed of small momenta. This is an extended description of the accumulating point discussed before in Ref. [13]. On the other hand, the solution (12) with $a_{\text{sn}} = 2NK(k)/L$ and $k \rightarrow 1$ needs large momentum components to describe its singular behavior at $x = 0$ and $\pm L$, which gives infinite harmonic resolution $M \rightarrow \infty$.

We would like to express sincere thanks to K. Yamawaki for valuable discussions. T.S. would like to thank K. Harada and K. Yazaki for helpful comments. T.S. was supported by the Grant-in-Aid for Scientific Research Fellowship, No. 11000480.

-
- [1] P.A.M. Dirac, Rev. Mod. Phys. **21**, 392 (1949).
 - [2] K.G. Wilson, T.S. Walhout, A. Harindranath, W. M. Zhang, R.J. Perry, and S.D. Glazek, Phys. Rev. D **49**, 6720 (1994); K. Yamawaki, “*New Nonperturbative Methods and Quantization on the Light Cone*”, Les Houches, France, Feb. 24-Mar. 7, 1997; S.J. Brodsky, H.C. Pauli, and S.S. Pinsky, Phys. Rept. **301**, 299-486 (1998).
 - [3] T. Banks, W. Fischler, S. H. Shenker, and L. Susskind, Phys. Rev. D **55**, 5112 (1997); L. Susskind, hep-th/9704080.
 - [4] H. Leutwyler, J.R. Klauder, and L. Streit, Nuovo Cim. **66A**, 536 (1970).
 - [5] T. Maskawa and K. Yamawaki, Prog. Theor. Phys. **56**, 270 (1976).
 - [6] A. Casher, Phys. Rev. D **14**, 452 (1976).
 - [7] H. C. Pauli and S. J. Brodsky, Phys. Rev. D **32**, 1993 (1985); *ibid* 2001 (1985); T. Eller, H.C. Pauli, and S.J. Brodsky, Phys. Rev. D **35**, 1493 (1987); T. Eller and H.C. Pauli, Z. Phys. C **42**, 59 (1989).
 - [8] Y. Kim, S. Tsujimaru, and K. Yamawaki, Phys. Rev. Lett. **74**, 4771 (1995); S. Tsujimaru and K. Yamawaki, Phys. Rev. D **57**, 4942 (1998).
 - [9] T. Heinzl, C. Stern, E. Werner and B. Zeller mann, Z. Phys. C **72**, 353 (1996); D.G. Robertson, Phys. Rev. D **47**, 2549 (1993); C.M. Bender, S.S. Pinsky, and B. van de Sande, Phys. Rev. D **48**, 816 (1993); S.S. Pinsky and B. van de Sande, Phys. Rev. D **49**, 2001 (1994); S.S. Pinsky, B. van de Sande, and J.R. Hiller, Phys. Rev. D **51**, 726 (1995).
 - [10] T. Sugihara, Phys. Rev. D **57**, 7373 (1998).
 - [11] J. S. Rozowsky and C. B. Thorn, Phys. Rev. Lett. **85**, 1614 (2000).
 - [12] S. Coleman, *Aspects of Symmetry* (Cambridge University Press, Cambridge, England, 1985).
 - [13] N. Nakanishi and K. Yamawaki, Nucl. Phys. B **122**, 15 (1977).